

## ***In Situ* Heating Studies of Sandia Octahedral Molecular Sieve**

A. Tripathi, Y. Lee, A. Celestian, J. Parise (SUNY, Stony Brook), and J.C. Hanson (BNL, NSLS)


Abstract No. Trip2954

Beamline(s): X7B

**Introduction:** SOMS-1 is a new octahedral molecular sieve developed at Sandia National Labs for selective removal of  $\text{Sr}^{2+}$  and its immobilization via phase transformation into a stable perovskite phase [1]. Based on these unique properties, it is an ideal candidate for  $^{90}\text{Sr}$  cleanup and sequestration at various DOE waste sites where the  $^{90}\text{Sr}$  is located in contaminated plumes and groundwater as mixed wastes. *In situ* dehydration studies were carried out using the as synthesized  $\text{Na}^+$  and acid ( $\text{H}^+$ ) exchanged phases. The H-exchanged phase transforms to a mixture of phases including an ilmenite phase whereas the as synthesized  $\text{Na}^+$  phase transforms to a cubic perovskite phase at ca. 500 °C (**Figure 1**). The structure of new hexagonal ilmenite phase is being studied by single crystal synchrotron X-ray diffraction studies at X3A1 beamline. The mechanism of ion-exchange will be determined via *in situ* ion-exchange studies during the next cycle.

**References:** [1] M.Nyman, A.Tripathi, J. Parise, R.S. Maxwell, W.T.A Harrison and T.M. Nenoff. (2000) Submitted in Nature

**Acknowledgments:** The authors thank the NSF for financial support (Grant DMR 97-13375). Research carried out in part at the NSLS at BNL is supported by the U.S. Department of Energy, Division of Materials Sciences and Division of Chemical Sciences, Office of Basic Energy Sciences.



**Figure 1.** Plot of the synchrotron X-ray powder diffraction profiles as a function of time during the 5-hr. dehydration of Na-SOMS-1